



Occurrence of Ectoparasites in Backyard Domestic Chickens (*Gallus gallus domesticus*) in the Northeast of Tunisia

KHALED KABOUDI^{1*}, RAFIKA BEN ROMDHANE², AMENI BEN SALEM³, MONCEF BOUZOUAIA¹

¹Department of Poultry Farming and Diseases, National Veterinary Medicine School of Sidi Thabet; University of Manouba, Tunisia; ²Department of Animal Production, Regional District of Agricultural development of Tunis – Tunisia; ³National Center of Zoosanitary Eve of Tunisia, Tunisia.

Abstract | The presence of ectoparasites on birds can lead to low productivity and eventually death. This study was carried out to identify and estimate the prevalence of ectoparasites of free-range poultry (*Gallus gallus*), in the Northeast of Tunisia. A total of 512 birds of both sexes and aged from 1.5 month to 24 months were examined and/or necropsied. Results showed that $19 \pm 3.39\%$ (98 birds), as overall prevalence, were infested by at least one parasite specie. Young animals were more infested ($13 \pm 2.91\%$) than adults ($6 \pm 2.05\%$). There was significant difference ($P < 0.05$) in prevalence found during rainy and cold seasons (winter: $24 \pm 6.59\%$; spring: $23 \pm 7.83\%$; autumn: $20 \pm 7.87\%$). Six species of ectoparasite were identified in the present study. *Cnemidocoptes mutans* ($8 \pm 2.35\%$) was the most prevalent, followed by *Menopon gallinae* ($6 \pm 2.05\%$), *Laminosioptes cysticola* ($3 \pm 1.47\%$), tick larvae of *Argas persicus* ($3 \pm 1.47\%$), *Echidnophaga gallinacea* ($3 \pm 1.47\%$) and *Menacanthus stramineus* ($2 \pm 1.22\%$). Infestation by one species of ectoparasite ($13 \pm 2.91\%$) was more frequently than mixed infestation ($6 \pm 2.05\%$). On the basis of these results it could be suggested that, good management, biosecurity and the education of farmers could be a vital measures to keep free from ectoparasites and improve the productivity of the chicken.

Keywords | Backyard poultry, Ectoparasite, Prevalence, Infestation, Tunisia

Editor | Asghar Ali Kamboh, Sindh Agriculture University, Tandojam, Pakistan.

Received | February 25, 2019; **Accepted** | June 18, 2019; **Published** | September 15, 2019

***Correspondence** | Khaled Kaboudi, Department of Poultry Farming and Diseases, National Veterinary Medicine School of Sidi Thabet; University of Manouba, Tunisia; **Email:** kaboudikhaled17@gmail.com

Citation | Kaboudi K, Romdhane RB, Salem AB, Bouzouaia M (2019). Occurrence of ectoparasites in backyard domestic chickens (*Gallus gallus domesticus*) in the northeast of Tunisia. J. Anim. Health Prod. 7(3): 92-98.

DOI | <http://dx.doi.org/10.17582/journal.jahp/2019/7.3.92.98>

ISSN | 2308-2801

Copyright © 2019 Kaboudi et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

INTRODUCTION

Backyard poultry sector is considered as an important source of income for several families in rural areas in Tunisia. These flocks provide a cheap source of protein in the form of meat and eggs. This traditional production contributes to the national poultry production with average of 7% (GIPAC, 2010).

Several factors limit the productivity in free-range poultry flocks. Malnutrition, poor management, lack of biosafety conditions and poor vaccination plans. Losses have also been attributed to limit housing and specific veterinary care services. Furthermore, poor genetic potential due to lack of selection and predation are also potential threats to

productivity. Parasitic infestation is considered as an important health problem in free-range poultry which handicaps production (Mungube et al., 2008).

Birds reared in the free range system are at constant risk of infestation by several types of endoparasites and ectoparasites. Several, species of ectoparasites (flies, lice, mites, and ticks) can infest poultry (Permin et al., 2002). They cause discomfort, decrease in growth and egg production, anaemia, irritation, loss of plumage and skin lesions that may be site of secondary infection. Death may occur when infestation is massive. Environments with high parasitic pressure can limit immunity protection against infectious diseases. A number of epidemiological factors including host, sex, age, breed and environment may influence the occurrence

Table 1: Prevalence rate of ectoparasites in male, female, young (≤ 18 weeks) and adult (> 18 weeks) free-range poultry

	Sex		χ^2 (p-value)	Age		χ^2 (p-value)
	Male	Female		Young	Adult	
Number of examined	151	361	1.02 (p=0.312)	382	130	4.389 (p=0.036)*
Number of positive	33	65		65	33	
Prevalence (%)	22 \pm 6.60%	18 \pm 3.96%		17 \pm 3.77%	25 \pm 7.44%	

* = significant difference

Table 2: Prevalence rate of ectoparasites in various types of free-range poultry

	Broiler	Pullet	Layer	Rooster	χ^2 (p-value)
Number of examined	138	244	117	13	57.24 (p<0.001)*
Number of positive	20	45	20	13	
Prevalence (%)	14 \pm 5.79%	18 \pm 4.82%	17 \pm 6.80%	100 \pm 23.07%	

* = significant difference

Table 3: Season-wise prevalence of ectoparasites in free-range poultry

Season	Number of examined	Number of positive	Prevalence (%)	χ^2 (p-value)
Autumn	99	20	20 \pm 7.87%	9.549 (p=0.022)*
Winter	161	38	24 \pm 6.59%	
Spring	111	25	23 \pm 7.83%	
Summer	141	15	11 \pm 5.16%	
Total	512	98	19 \pm 3.39%	

* = significant difference

and intensity of parasitic infestations (Nadeem et al., 2007).

Ectoparasites feed on blood, feathers, skin and scales of their host. They may transmit several infectious diseases and serve as transport or intermediate hosts for different helminthic parasites (Hopla et al., 1994). *Menacanthus stramineus*, known to feed on blood, can carry the equine encephalomyelitis virus. *Chlamydophila psittaci*, has been isolated from *Menopon gallinae* (Calnek et al., 1991). Many other poultry pathogens, such as *Pasteurella*, fowl pox, Newcastle disease virus, can be spread by ticks and mites (Nnadi and George, 2010). *Dermanyssus gallinae* has been widely reported to transmit human and animal pathogens (e.g., viruses and bacteria) and parasites (e.g., Hepatozoon) to persons (Valiente et al., 2009).

The present study was undertaken with the aims of determining the prevalence rates of ectoparasitic species in free-range backyard chickens in the Northeast regions of Tunisia.

MATERIALS AND METHODS

This study was conducted, from September 2011 to June 2015, on 512 local chickens. Animals are from regions rounding the town of Sidi Thabet, in the Northeast of Tunisia. All birds were from farms in which the traditional breeding is practiced. They were of both sexes and aged

from 1.5 to 24 months. These animals were divided into two groups, namely young and adult.

The birds were received for clinical exam and/or necropsy in the Avian Clinic of the National Veterinary Medicine School of Tunisia. External parasites were collected from different parts of the body around the wing, head, vent, feather, feet, leg wattle and comb. The legs and featureless areas with crest were scrapped for parasitological identification. Parasites were identified according to their morphological characteristics using the key criteria as mentioned by Soulsby (1982).

STATISTICAL ANALYSIS

Comparative analysis of prevalence in chickens was performed using the chi-square test (Word Excel, Microsoft office, version 2013). A p-value less than or equal to 0.05 were used as a limit of statistical significance.

RESULTS

Overall, 98 (19 \pm 3.39%) of the examined birds were found to be infested with at least one ectoparasite specie. The infestation was predominated in male (22 \pm 6.60%) ($\chi^2 = 1.02$; p=0.312). The infestation was significantly higher in adult birds (25 \pm 7.44%) than in young (17 \pm 3.77%) ($\chi^2 = 4.389$; p=0.036) (Table 1).

Table 4: Prevalence of individual parasites in female, male, young (≤ 18 weeks) and adult (> 18 weeks) free-range poultry

Parasite species	Overall prevalence	No. positive (%)		χ^2 (p-value)	No. positive (%)		χ^2 (p-value)
		Female	Male		Young	Adult	
<i>Cnemidoptes mutans</i>	8 \pm 2.35%	12 (12 \pm 6.34%)	31 (32 \pm 9.23%)	50.636 (p<0.001)*	18 (18 \pm 7.6%)	25 (26 \pm 8.57%)	20.535 (p<0.001)*
<i>Menopon gallinae</i>	6 \pm 2.05%	15 (15 \pm 7.07%)	14 (14 \pm 6.87%)	3.932 (p=0.02)*	20 (20 \pm 7.92%)	9 (9 \pm 5.66%)	0.128 (p=0.5)
<i>Menacanthus stramineus</i>	2 \pm 1.22%	5 (5 \pm 4.31%)	7 (7 \pm 5.05%)	3.723 (p=0.05)*	4 (4 \pm 3.88%)	8 (8 \pm 5.37%)	6.664 (p=0.001)*
<i>Echidnophaga gallinacea</i>	3 \pm 1.47%	4 (4 \pm 3.88%)	10 (10 \pm 5.94%)	10.424 (p=0.001)*	11 (11 \pm 6.19%)	3 (3 \pm 3.37%)	1.096 (p=0.2)
<i>Laminosioptes cysticola</i>	3 \pm 1.47%	10 (10 \pm 5.94%)	7 (7 \pm 5.05%)	0.518 (p=0.3)	9 (9 \pm 5.66%)	8 (8 \pm 5.37%)	1.65 (p=0.1)
<i>Argas persicus</i>	3 \pm 1.47 %	8 (8 \pm 5.37%)	9 (9 \pm 5.66%)	3.419 (p=0.05)*	6 (6 \pm 4.7%)	11 (11 \pm 6.19%)	8.868 (p=0.001)*

* = significant difference

Different types of poultry were infested as indicated in Table 2. However, it was remarkable that all roosters were positive (100%), followed by pullets (18 \pm 4.82%), layers (17 \pm 6.80%) and broilers (14 \pm 5.79%). The relationship between infestation prevalence and type of production was highly significant ($\chi^2=57.24$; p<0.001).

The study according the season is summarized in Table 3. The results showed that parasitic infestations were diagnosed along the year, with statistically significant ($\chi^2=9.549$; p=0.022) most high incidence observed during winter (24 \pm 6.59%), spring (23 \pm 7.83%) and autumn (20 \pm 7.87%). Influence of sex ($\chi^2=9.495$; p=0.02), age ($\chi^2=14.713$; p=0.001) and type of production ($\chi^2=29.566$; p<0.001) on seasonal repartition was statistically significant.

Regarding to the importance of infestation, highest prevalence of single infestation (67 birds; 13 \pm 2.91%) was revealed compared to mixed infestation (31 birds; 6 \pm 2.05%). This repartition was influenced by the age ($\chi^2=10.149$; p<0.05) of birds but not by the sex factor ($\chi^2=0.515$; p=0.47). Study of mixed infestation showed a prevalence rate of 5 \pm 1.88% (28 birds) and 1 \pm 0.86% (3 birds) in chickens infested by two and by three species of ectoparasites, respectively. No animals infested by up than three ectoparasites in this study.

Six ectoparasite species were identified in the present study. Results of the individual prevalence of each parasite depending on sex and age were summarized in Table 4. The most prevalent species was *Cnemidoptes mutans* (8 \pm 2.35%) (Figure 1), followed by *Menopon gallinae* (6 \pm 2.05%) (Figure 2), tick larvae of *Argas persicus* (Figure 3), *Laminosioptes cysticola* (Figure 4) and *Echidnophaga gallinacea* with the same percentage (3 \pm 1.47%). Whereas, *Menacanthus stramineus* (2 \pm 1.22%) (Figure 5) was the least prevalent ectoparasite, identified in backyard chickens in

the studied region.



Figure 1: *Cnemidoptes mutans* in free-range chicken



Figure 2: *Menopon gallinae* in free-range chicken

DISCUSSION



Figure 3: Severe infestation of a free-range chicken by larvae of *Argas persicus*. (Note hemorrhage zones corresponding to the points of attachment of tick larvae)



Figure 4: Calcified cysts of *Laminosioptes cysticola* (arrows) in the subcutaneous collagen in the region of breast



Figure 5: *Menacanthus stramineus* in free-range chicken

Results of this study show overall prevalence assessed to $19 \pm 3.39\%$ (98 animals out of 512 examined) in the north-east region of Tunisia. High prevalence was found in other countries, such as Zimbabwe (Goromonzi District) (100%) (Permin et al., 2002), Potiskum (Nigeria) (84.50%) (Lawal et al., 2017), Benue State (Nigeria) (69.7%) (Oche et al., 2016), West of Iran (64.83%) (Rezaei et al., 2016) and Algeria (69-100%, according to the region study) (Meguini et al., 2018). Infestation by ectoparasites is favored in free range farming system by several factors, such as possible contact of chickens with animals of other poultry species and wild birds (Adelusi et al., 2014).

Prevalence of the ectoparasites infestation was found to be statistically higher in adult ($25 \pm 7.44\%$) than in young. Malann et al. (2016), Oche et al. (2016), Kebede et al. (2017) and Lawal et al. (2017) have also reported that adult village chickens were more infested by ectoparasites compared with younger ones. This finding might be associated with the frequent contact of adult chickens with other species of animals and may be exposed longer to the infested environment and other source of infestation. These findings were in coherence with prevalence rate calculated in different type of production. In fact, roosters (100%) and pellets ($18 \pm 4.82\%$) were statistical significantly more frequently infested by ectoparasites than layers ($17 \pm 6.80\%$) and broilers ($14 \pm 5.79\%$).

Our results regarding the age of animal were in disagreement with the findings of Mulugeta et al. (2013), Firaol et al. (2014) and Rezaei et al. (2016) who reported that young chickens were more infested than adult animals.

There is no affinity of ectoparasites to the sex of animals, which is in disagreement with the findings of Tolossa et al. (2009). Indeed, these authors demonstrated higher significance infestation of male (93.94%) compared to female (76.19%). Prevalence of ectoparasitic infestation was statistical significantly higher in raining and humid season (autumn, spring and winter), with global prevalence assessed to $22 \pm 4.21\%$, which is in agreement with the observations of Mukaratirwa and Hove (2009) and Mohammad et al. (2016) who also reported high prevalence of ectoparasites in chickens during the rainy season. Generally, the rainy season consists of favorable climatic conditions for the proliferation of parasites. Moderated to high ambient temperature and humidity during this season are very essential for the hatching of eggs and larval developmental stages (Soulsby, 1982). These climatic characteristics explain that external parasites of poultry are common in the tropics (Nnadi and George, 2010; Firaol et al., 2014). Although, ectoparasites infestation have also been reported

during the dry season (Alemu et al., 2015; Kebede et al., 2017).

Single ectoparasite infestation ($13 \pm 2.91\%$) found in this study was more frequent than mixed infestation ($6 \pm 2.05\%$). These findings are not in line with those of several authors, such as Medjouel et al. (2013), Mulugeta et al. (2013), Firaol et al. (2014) and Lawal et al. (2017), who reported higher prevalence of mixed ectoparasitic infection in village chickens. The difference may be associated with different agro-ecology, climatic factors in the various study areas, variation in sample size and sampling period, free range nature of village chickens, level of hygiene in and around the poultry houses and the level of parasite control practices.

Six species of ectoparasites were identified in the present study, which included *Cnemidocoptes mutans*, *Menopon gallinae*, *Menacanthus stramineus*, *Echidnophaga gallinacea*, *Laminosioptes cysticola* and tick larvae of *Argas persicus*. Among these species of ectoparasites encountered, *Cnemidocoptes mutans* ($8 \pm 2.35\%$) was the most prevalent, while *Menacanthus stramineus* ($2 \pm 1.22\%$) was the least. The prevalence rate of *Cnemidocoptes mutans* was higher than that mentioned by Lawal et al. (2017) (7%) and Odeno et al. (2016) (7.26%). Nevertheless, it was lower than prevalence rate found by Bala et al. (2011) (9.40%), Firaol et al. (2014) (34.62%) and Oche et al. (2016) (18.18%). *Cnemidocoptes mutans* is a scale leg mite; a small spherical sarcoptic mite usually tunnels in to the tissue under the scales causing an inflammation and market keratinization, which is responsible for the thickened scaly nature of the feet. Low prevalence revealed in our study compared to the prevalence reported by Hagos and Eshetu (2005) (19.5%), may be due to the cold temperature and/or health management system.

Among the lice species observed, *Menopon gallinae* and *Menacanthus stramineus*. The prevalence rate of *Menopon gallinae* from the present study was $6 \pm 2.05\%$, which were considered higher than that found by Lawal et al. (2017) in Nigeria. However, our result was lower than 13.28%, 8.1%, 12.4%, 40.12%, 97.2% and 97.7%, reported by Kebede et al. (2017), Bala et al. (2011), Moyo et al. (2015), Sadiq et al. (2003), Medjouel et al. (2013) and Sabuni et al. (2010), respectively. *Menacanthus stramineus* was detected in $2 \pm 1.22\%$ of examined chickens. Its prevalence rate was lower than results reported by several authors in different countries, such as Belihu et al. (2010), Bala et al. (2011), Medjouel et al. (2013), Moyo et al. (2015) and Mirullo and Mullens (2016) in Ethiopia, Nigeria, Algeria, South Africa and California, respectively. This parasite is the most pathogenic in poultry. It causes severe anemia in the host by feeding on blood that oozes out. It also causes inflammation of skin and extensive scab formation. However, exact

impact of lice species on the chicken was not determined. The prevalence rate of *Laminosioptes cysticola* was $3 \pm 1.47\%$. This ectoparasite was rarely described in published studies. Considered as a parasite of the subcutaneous and inter-muscular collagen tissue, it is generally not pathogenic in poultry. However, this parasite may represent a risk of food-borne human allergy. Infestation induce the formation of white calcified nodules observed in the necropsy exam or after slaughter in the internal face of thigh, on the breast and the neck and on the internal viscera (Da Silva Martins et al. 2010). Sometimes, severe infestation can induce clinical illness. In this way, Smith et al. (1997) described nervous signs (torticollis, circling, loss of balance, wing droop) in wild turkey massively infested with *Laminosioptes cysticola*. Indeed, granulomatous inflammation can be abundant in brachial plexus and sciatic nerves.

Tick larvae infestation were represented by *Argas persicus* in $3 \pm 1.47\%$ of cases. This prevalence was lower than that mentioned by Lawal et al. (2017) (6.2%) and Rezaei et al. (2016) (78.66%) in Nigeria and Iran, respectively. High risk of infestation was related to the presence of other animal species (cattle, dog...) in the farm. In this condition, ticks causes blood spoliation leading anemia and death. *Argas persicus* can cause paralysis in infested birds and a decrease in egg production. Indirect pathogenic action of ticks is explained by their role of pathogens vector. *Argas* spp can infest man, especially in rural areas, where there close contact between man and animals. They cause irritation and allergic problems (Sadiq et al., 2009). In fact, these parasites can transmit bacterial, rickettsial, viral, parasitic and spirochaetal diseases in poultry (Haider Shah et al., 2004). In the same way, *Argas persicus* larvae have been responsible for episodes of infectious bursal disease and spirochaetosis (Abdu, 1987).

Among the identified flea species, *Echidnophaga gallinacea* was detected in $2 \pm 1.22\%$ of examined birds. The prevalence of this ectoparasite in our study was not similar to those reported from west of Iran with 6% (Rezaei et al., 2016), 18% in California (Murillo and Mullens, 2016), 29.6% in Abuja (Nigeria) (Odeno et al., 2016), 27.3% in Gombe (Nigeria) (Lawal et al., 2016) and from different regions in Ethiopia with 26.6% (Mata et al., 2018) to 51.3% (Belihu et al., 2009). *Echidnophaga gallinacea* attaches to the combs, wattles and around the eyelids, where it induces irritation, restlessness and anemia (Soulsby, 1982). Our study excludes many other parasites feeding only briefly on birds, such as *Goniocotes gallinae*, *Lipeurus caponis* and *Dermanyssus gallinae*.

Finally, it is important to mentioning that variations in the prevalence rates and diversity of species of ectoparasites from the reports of the various studies may be connect to several factors. The differences in geographical are-

as, host factors, husbandry and management system, poor sanitation, sample size, period of study as well as different favorable climatic conditions such as temperature and humidity which may influence the population dynamics of the ectoparasites are the most included epidemiological factors (Arends, 2003; Prelezov and Kolnarski, 2006).

CONCLUSION

This study showed that backyard chicken could be infested by several species of ectoparasites in the regions around Sidi Thabet in the Northeast of Tunisia. Parasitic infestation affect health status of animals and their productivity. Several risk factors can influence the prevalence and the severity of infestation, such as age, sex and season. Poor management conditions, mixed animal and avian species and negligence of antiparasitic application can preserve the infestation in the flock. That is why, the association of insecticide treatments to the hygiene are crucial to minimize damages and preserve the traditional poultry. Further studies are needed to determine the economic impact of ectoparasitic infestation in backyard flocks and to perform global strategy of control.

CONFLICT OF INTEREST

There is no conflict of interest.

AUTHORS CONTRIBUTION

Khaled Kaboudi: Sampling, statistical analysis, interpretation and redaction.

Rafika Ben Romdhane: Sampling and redaction.

Ameni Ben Salem: Sampling and interpretation.

Moncef Bouzouaia: Interpretation.

REFERENCES

- Abdu PA (1987). Infectious bursal disease in pullet chicks. Avian Dis. 31: 204-205.
- Adelusi SM, Vajime CG, Omudu EA, Okpotu RO, Onazi FO (2014). Avian ectoparasitism in Makurdi, Nigeria: Do wild birds serve as reservoir for domestic birds? Nigerian J. Pure Appl. Sci. 6: 11-15.
- Alemu N, Muktar Y, Kassaye D Hiko A (2015). Prevalence of lice and fleas in backyard chickens of Bishoftu Town, Ethiopia. American-Eurasian J. Agric. Environ. Sci. 15 (11): 2136-2142.
- Arends JJ (2003). External parasites and poultry pests. In: Diseases of poultry; 11 ed., (Edited by Calnek WB, Barnes JH, Beard WC, Mc Dougald LR, Saif YM. Iowa State Press, Blackwell Publishing Company, Ames, Iowa, Pp: 905-930.
- Bala AY, Anka SA, Waziri A Shehu H (2011). Preliminary Survey of Ectoparasites Infesting Chickens (*Gallus domesticus*) in Four Areas of Sokoto Metropolis. Nig. J. Basic App. Sci. 19: 173-180.
- Belihu K, Mamo A, Lobago F, Ayana D (2010). Prevalence of ectoparasites in backyard local chickens in three agroecologic zones of East Shoa in Ethiopia. Revue Méd. Vét. 160: 537-541.
- Calnek BW, Barnes HJ, Bread CW, Reid WM, Yoder HW (1991). Diseases of Poultry. 9th ed. Iowa: Iowa State University Press.
- Da Silva Martins NR, de Resende JS, Marques MVR, Barrios PR, Fernandes AA, Montresor LC, Cunha LM (2010). *Laminosioptes cysticola* in free-range chickens in Minas Gerais, Brazil. Ciência Rural, Santa Maria. 40(6): 460-463.
- Firaol T, Dagmawit A, Askale G, Solomon S, Morka D, Waktole T (2014). Prevalence of ectoparasite infestation in chicken in and around Ambo Town, Ethiopia. J. Vet. Sci. Technol. 5: 189. <https://doi.org/10.4172/2157-7579.1000189>
- GIPAC (2010). Groupement Interprofessionnel des Produits Avicoles et Cunicoles. (www.gipaweb.com.tn)
- Hagos A, Eshetu Y (2005). Ectoparasites of focal scavenging chickens of central Ethiopia SINET. Ethiopia J. Sci. 28(1): 69-74.
- Haider Shah A, Nisar Khan M, Iqbal Z, Sohail Sajid M (2004). Tick infestation in poultry. Int. J. Agric. Biolog. Engineering 6:1162-1165.
- Hopla CE, Durden LA, Keirans JE (1994). Ectoparasites and classification. Rev. Sci. Tech. 13: 985-1017.
- Kebede A, Abebe B, Zewdie T (2017). Study on prevalence of ectoparasites of poultry in and around Jimma Town. Eur. J. Biol. Sci. 9 (1): 18-26. <https://doi.org/10.5829/idosi.ejbs.2017.18.26>
- Lawal JR, Yusuf ZB, Dauda J, Gazali YA, Biu AA (2017). Ectoparasites infestation and its associated risk factors in village chickens (*Gallus gallus domesticus*) in and around Potiskum, Yobe State, Nigeria. J. Anim. Husb. Dairy Sci. 1(1): 8-19.
- Malann D, Olatunji O, Usman M (2016). Ectoparasitic infestation on poultry birds raised in Gwagwalada area council, FCT-Abuja. Int. J. Innov. Res. Develop. 5(13): 74-77.
- Mata W, Galgalo W, Jilo K (2018). Prevalence of the major ectoparasites of poultry in extensive and intensive farms in Jimma, Southwestern Ethiopia. J. Parasitol. Vector Biol. 10(7): 87-96. <https://doi.org/10.5897/JPV2017.0298>
- Medjouel I, Benakhla A, Senouci K, Djelil H, Matallah F (2013). Prevalence and distribution of chewing lice (Phthiraptera) in free range chickens from the traditional rearing system in the Algerian North East, Area of El-Tarf. Int. J. Poult. Sci. 12: 721-725. <https://doi.org/10.3923/ijps.2013.721.725>
- Meguini MN, Righi S, Zeroual F, Saidani K, Benakhla A (2018). Inventory of lice of mammals and farmyard chicken in North-eastern Algeria. Vet. World 11(3): 386-396. <https://doi.org/10.14202/vetworld.2018.386-396>
- Mirullo AC, Mullens BA (2016). Diversity and prevalence of ectoparasites on backyard chicken flocks in California. J. Med. Entomol. 1-5. <https://doi.org/10.1093/jme/tjv243>
- Mohammad M, Omid G, Mohammad Y (2016). Prevalence of Ectoparasites of indigenous chickens from Dalahu Region Kermanshah Province, Iran. Turkiye Parazitolo. Derg. 40: 13-6. <https://doi.org/10.5152/tpd.2016.4185>
- Moyo S, Masika P J, Moyo B (2015). A Diagnostic Survey of External Parasites of Free-Range chickens, in the rural areas of Eastern Cape, South Africa. Int. J. Agric. Sci. Vet. Med. 3(2): 1 – 9.
- Mukaratirwa S, Hove T (2009). A survey of ectoparasites,

- cestodes and management of free-range indigenous chickens in rural Zimbabwe. *J. South African Vet. Assoc.* 80 (3): 188-191.
- Mulugeta A, Mersha C, Basaznew B (2013). Major constraints of village poultry production in Demba Gofa District of southern region, Ethiopia. *Br. J. Poult. Sci.* 2 (1): 01-06.
 - Mungube EO, Baun SM, Tenhagen BA, Wamae LW, Nzioka SM, Muhammed L, Nginyi JM (2008). Prevalence of Parasites of the Local Scavenging Chickens in a Selected Semi-Arid Zone of Eastern Kenya. *Trop. Anim. Health Prod.* 40: 101-109. <https://doi.org/10.1007/s11250-007-9068-3>
 - Nadeem M, Khan MN, Iqbal Z, Sajid MS, Arshad M, Yaseen M (2007). Determinants influencing prevalence of louse infestations of layers of district Faisalabad (Pakistan). *Br. Poult. Sci.* 485: 546-550. <https://doi.org/10.1080/00071660701573086>
 - Nnadi PA, George SO (2010). A cross-sectional survey on parasites of chickens in selected villages in the subhumid zones of south-Eastern Nigeria. *J. Parasitol. Res.* Article ID 141824, 6 pages. <https://doi.org/10.1155/2010/141824>
 - Oche DA, Ogwiji O, Torhemen M, Andyar AC, Akahaan RT, Luga ZT, Wayo NS, Tiough SM, Kwaghtse WR (2016). Survey of ecto-parasites in chickens in Benue State of Nigeria. *Scholarly J. Agricult. Sci.* 6(8): 235-238.
 - Odeno RA, Mohammed BR, Simon MK, Agbede RIS (2016). Ecto-parasites of Domestic Chickens (*Gallus gallus domesticus*) in Gwagwalada Area Council, Abuja, Nigeria-West Africa. *Alex. J. Vet. Sci.* 51(1): 140-146. <https://doi.org/10.5455/ajvs.220654>
 - Permin A, Esmann JB, Hoj CH, Hove T, Mukaratirwa S (2002). Ecto-, endo- and haemoparasites in the free-range chickens in the Goromonzi district in Zimbabwe. *Prev. Vet. Med.* 54: 213-224. [https://doi.org/10.1016/S0167-5877\(02\)00024-7](https://doi.org/10.1016/S0167-5877(02)00024-7)
 - Prelezov PN, Koinarski K T (2006). Species Variety and Population Structure of Mallophaga (Insecta: phthiraptera) on Chickens in the Region of Stara Zagora". *Bulgarian J. Vet. Med.* 9 (3): 193-200.
 - Rezaei F, Hashemnia M, Chalechale A, Seidi S Gholizadeh S (2016). Prevalence of ectoparasites in free-range backyard chickens, domestic pigeons (*Columba livia domestica*) and turkeys of Kermanshah province, west of Iran. *J. Parasit. Dis.* 40(2): 448-453. <https://doi.org/10.1007/s12639-014-0524-5>
 - Sabuni ZA, Mbuthia PG, Maingi N, Nyaga PN, Njagi LW, Bebora LC, Michieka JN (2010). Prevalence of ectoparasites infestation in indigenous free-ranging village chickens in different agro-ecological zones in Kenya. *Livestock Res. Rural Develop.* 22 (11): 1-5.
 - Sadiq NA, Adejinmi JO, Adedokun OA, Fashanu SO, Alimi AA, Sofunmade YT (2003). Ectoparasites and haemoparasites of indigenous chicken (*Gallus domesticus*) in Ibadan and environs. *Trop. Vet.* 21:187-191. <http://dx.doi.org/10.4314/tv.v21i4.4541>
 - Smith KE, Quist CF, Crum JM (1997). Clinical illness in a wild turkey with *Laminosioptes cysticola* infestation of the viscera and peripheral nerves. *Avian Dis.* 41: 484-489.
 - Soulsby E JL (1982). Helminths, arthropods and protozoa of domestic animals. 7th ed. London, UK: Bailliere and Tindal, 809 pages.
 - Tolossa YH, Shafi ZD, Basu AK (2009). Ectoparasites and gastrointestinal helminths of chickens of three agro-climatic zones in Oromia region, Ethiopia. *Anim. Biol.* 59: 289-297. <https://doi.org/10.1163/157075609X454926>
 - Valiente Moro C, De Luna CJ, Tod A, Guy JH, Sparagano OA, Zenner L (2009). The poultry red mite (*Dermanyssus gallinae*): a potential vector of pathogenic agents. *Experim. Appl. Acarol.* 48: 93-104.