

## Research Article



# Impact of Stocking Density on Growth Performance, Carcass Traits, and Economic Feasibility of Growing Rabbits

MOHAMED A.E. OMAR, MARWA EL-SHAHAT, FARDOS A.M. HASSAN\*

*Department of Animal Wealth Development, Faculty of Veterinary Medicine, Zagazig University, El-Zeraa St. 114; 44511-Zagazig, Egypt.*

**Abstract** | A total of sixty, 4 week-old male New Zealand White (NZW) rabbits were used to explore the impact of stocking density on the performance, carcass characteristics, blood biomarkers, and economic indices in growing rabbits. All animals were kept in wire cages (50 length x 50 width x 40 cm high) in groups of 2, 4, and 6 rabbits/cage (represented 8, 16, and 24 rabbits/m<sup>2</sup>); each stocking density was replicated 5 times. A significantly lower body weight (1748.67, 1567.58 vs. 2017.83 g) and feed intake (5040.36, 4807.47 vs. 5107.41 g) with higher feed conversion (4.22, 4.77 vs. 3.53) were observed in rabbits stocked 16 and 24 rabbits/m<sup>2</sup> compared to 8 rabbits/m<sup>2</sup>, respectively ( $P < 0.01$ ). No statistical variation ( $P > 0.05$ ) was found in percentages of skin, full stomach, full intestine, head, and carcass parts under different stocking densities. However, a significantly lower dressing out and liver percentage (52.42 and 5.12 %, respectively) was recorded in rabbits stocked at 24 rabbits/m<sup>2</sup> than other groups ( $P < 0.001$ ). Rabbits stocked at 24 rabbits/m<sup>2</sup> had the lowest values of RBCs counts, Hb, and PCV ( $P < 0.01$ ). The levels of total protein, globulin, urea, and creatinine were significantly lower in the group of 24 rabbits/m<sup>2</sup> than those of 8 and 14 rabbits/m<sup>2</sup> ( $P < 0.05$ ). The total and net revenue was significantly decreased and the cost-benefit ratio was increased from 0.61 to 0.77 as the number of rabbits increased from 8 to 24 rabbits/m<sup>2</sup> ( $P < 0.001$ ). It can be concluded that high stocking density adversely affects rabbit performance and profitability.

**Keywords** | Performance, Carcass traits, Cost analysis, Rabbits

**Editor** | Asghar Ali Kamboh, Sindh Agriculture University, Tandojam, Pakistan.

**Received** | November 02, 2020; **Accepted** | November 23, 2020; **Published** | December 27, 2020

\***Correspondence** | Fardos A.M. Hassan, Department of Animal Wealth Development, Faculty of Veterinary Medicine, Zagazig University, El-Zeraa St. 114; 44511-Zagazig, Egypt; **Email:** fardoseconomy@yahoo.com

**Citation** | Omar MAE, El-Shahat M, Hassan FAM (2020). Impact of stocking density on growth performance, carcass traits, and economic feasibility of growing rabbits. *J. Anim. Health Prod.* 9(s1): 50-55.

**DOI** | <http://dx.doi.org/10.17582/journal.jahp/2020/9.s1.50.55>

**ISSN** | 2308-2801

**Copyright** © 2020 Omar *et al.* This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

## INTRODUCTION

The rabbit production has recently gained popularity due to its high productivity, fast growth rate, efficient feed conversion ratio, and a better return on capital which are tantamount to that of broiler chicken (Gondret *et al.*, 2005). Rabbits can be fed a diet containing up to 30% crude fiber as opposed to 10% by most poultry species and can change 20% of the protein they consume into palatable meat (Egbo *et al.*, 2001). Rabbit meat is of high quality protein with lower calories, fat, and cholesterol levels relative to other sources of meat (Samkol and Lukefahr, 2008).

In the rabbit sub-sector, farmers must focus on optimizing kit longevity and growth efficiency; one of the best approaches to accomplish this is to establish a conducive raising environment with an appropriate stocking rate (El-Bayoumi *et al.*, 2018). Indeed, stocking density can be defined as the number of animals per square meter in a cage, pen, or building. It is a crucial factor that can influence labor cost, investment cost, performance, and thus the profitability of rabbit production (Grace and Olorunju, 2005). The stocking densities and minimum space allowance for rabbits should often apply to the final weight achieved by animals (of a certain strain, sex, and feeding regime). The European Food and Safety Authority

(2005) suggested a base space of 625cm<sup>2</sup>/rabbit; not more than 40 kg/m<sup>2</sup> at the end of fattening to discourage rabbit behavior disorder. In particular, the ideal cage space should allow each growing rabbit to stretch along one side of the cage and sit upright at all intervals of age (Onbaslar and Onbaslar, 2007).

The economic value of stocking density derives from its negative impact on gain per animal housed and obtaining too small profit margins as overcrowding limits the space available for eating and drinking and raises aggressive animal actions (Bigler and Oester, 1996). In overcrowded cages, the nutritional and locomotive behavior of rabbits changes, which affects their carcass characteristics and meat quality (Szendrő and Dalle Zotte, 2011). In this regard, it is pertinent to explore the harmony between farm returns and a sensible degree of animal welfare regarding stocking density. Therefore, this research aimed to investigate the influence of different stocking density on growth performance, carcass characteristics, blood biomarkers, and economic efficiency of New Zealand White (NZW) rabbits.

## MATERIALS AND METHODS

The experiment was conducted in compliance with the ethical standards and regulations set out by the Faculty of Veterinary Medicine’s Local Laboratory Animals Care Committee, Zagazig University. The experimental practice had been approved by the institutional ethics committee (Approval No. ANWD 206).

### ANIMALS, MANAGEMENT AND EXPERIMENTAL DESIGN

A total of sixty 4 week-old male NZW rabbits (561.7±8.3 g) obtained from a commercial rabbit farm, and were utilized in this experiment. Rabbits were randomly assigned into three groups of 10, 20, and 30 rabbits, and each group was subdivided into five replicate cages. Each replicate cage in the groups comprised of 2, 4, and 6 rabbits that correspond 8, 16, and 24 rabbits/m<sup>2</sup>, respectively; where each cage measured 50 x 50 x 40 cm (length x width x high). Throughout the experiment, the lighting schedule was kept at 16 h of light and 8 h of darkness; the surrounding temperature was adjusted at 23 ±2°C and the relative humidity was about 65 ±5%. All diets were given in the pelleted form, and the basal diet was designed to fulfill the nutritional needs as per NRC Recommendations (1977; Table 1). The animals had an *ad libitum* excess to feed and water. The experimental procedures continued for 7 consecutive weeks (from week 4 to week 11 of age).

### PERFORMANCE TRAITS

Rabbits’ body weights were measured at the beginning (week 4; initial weight) and the end of the experimental

trial (week 11; final weight), and body weight gain (BWG) was estimated as the difference between final weight and initial weight. Cage feed intake (FI) was measured over this period, and the feed conversion rate (FCR) was calculated as the ratio of feed (g) to weight gain (g). The mortality rate was recorded daily throughout the experimental period. The rabbits died during the experiment were weighed, and all feed losses in each cage were gathered and weighed to ensure FI and FCR accuracy.

**Table 1:** Composition and chemical analyses of the basal diet (as fed basis).

Ingredients (%)	Basal diet
Yellow corn	15.3
Soybean meal, 44% CP	19.0
Wheat bran	14.0
Alfalfahay, 12% CP	32.2
Barley	12.0
Corn gluten	1.0
Molasses	3.0
Dicalcium phosphate	0.7
Sodium chloride	0.3
Vit. and min. premix*	0.3
Limestone	1.0
DL-Methionine, 98%	0.1
Toxinil	0.1
<b>Analysed nutrient content (%)†</b>	
Crude protein	17.4
Ether extract	2.2
Crude fibre	14.3
Crude ash	7.6
Dry matter	89.2
Organic matter	92.4
Digestible energy, kcal/kg	2508.9

\* Vitamin and Mineral premix, provided per kg of diet: Vitamin A, 6000 IU; Vitamin D3, 1200 IU; Vitamin E, 26 IU; Vitamin K3, 1 mg; Vitamin B1, 1 mg; Vitamin B2, 3 mg; Vitamin B6, 1.5 mg; Pantothenic acid, 10 mg; Vitamin B12, 3.5 µg; Niacin, 30 mg; Folic acid, 1.25 mg; Biotin, 16 µg; Fe, 50 mg; Zn, 50 mg; Mn, 20 mg; Cu, 6 mg; I, 0.3 mg; Se, 0.03 mg; Choline chloride, 500 mg. † Analysis was performed according to AOAC (2000).

### CARCASS TRAITS

At the end of the study, five rabbits per group with average body weight for the corresponding group were chosen for blood sampling and carcass evaluation, earlier which they fasted for 12 h. All rabbits were weighed (representing slaughter weight, SW), euthanized by cutting the two jugulars, and then the skin, gastrointestinal tract, and distal leg segment were detached. Eviscerated carcasses (with head, liver, kidneys, thoracic cage organs, perirenal fat) were ice-cold for 24 h at 4°C and then weighed to get the chilled

carcasses weight (CCW). After that, the head, thoracic organs, liver, and kidneys were excised and the carcasses were weighed again to obtain the reference carcasses weight (RCW). The dressing out (CCW as a percentage of SW) and the proportion of head, liver, kidney, and carcass parts (fore, mid, and hind parts) to either CCW or RCW were evaluated as required.

### HEMATOLOGICAL AND SERUM BIOCHEMICAL PARAMETERS

Blood samples (n= 5/group) were taken during the slaughter process, one part was transferred to a vacutainer tube containing EDTA as an anticoagulant for hematological evaluation and the other part was transferred to a gel activator tube for serum separation (3000 rpm; 15 min; 4 °C). Red blood cells count (RBC's), white blood cells count (WBC's), hemoglobin concentration (Hb), packed cell volume (PCV), mean corpuscular volume (MCV), and mean cell hemoglobin (MCH) of the whole blood specimens were measured using a Hema Screen 18 automated hematology analyzer (Hospitex Diagnostics, Sesto Fiorentino, Italy), whereas differential leukocyte counts (lymphocytes and heterophils) were enumerated manually according to Dacie and Lewis (1991). The heterophil to lymphocyte (H:L) ratio was also determined.

Serum total protein, albumin, cholesterol, triglycerides, alanine aminotransferase (ALT), aspartate aminotransferase (AST), creatinine and urea were investigated using commercial diagnostic kits (Diamond Diagnostics, Holliston, MA, USA) according to the producer's guidelines.

### ECONOMIC FEASIBILITY MEASUREMENTS

A partial budgeting technique was utilized to evaluate the economic impact of stocking density as follow:

$$\text{Total feed cost} = \text{Total feed intake per rabbit} \times \text{cost of one kg diet}$$

Total cost was estimated by considering feeding cost as well as the expense of experimental rabbits, veterinary services, labor and other miscellaneous expenses.

$$\text{Total revenue} = \text{rabbit live body weight} \times \text{price of kg live BW.}$$

$$\text{Net revenue} = \text{total revenue} - \text{total cost}$$

$$\text{Cost-benefit ratio} = \text{Total cost/total revenue}$$

### STATISTICAL ANALYSIS

Data were evaluated by one-way analysis of variance (ANOVA) using the GLM procedure in SPSS (SPSS Inc., Chicago, Illinois, USA), with individual animal or cage replicate served as the experimental unit for all statistical analyses. Shapiro-Wilk's test was applied to verify the normal distribution of data, and Levene's test to ascertain the homogeneity of variance components between experimental groups. The significance of the differences

between the mean values was estimated using Bonferroni test, where  $P < 0.05$  was considered significant.

## RESULTS AND DISCUSSION

### IMPACT OF STOCKING DENSITY ON PERFORMANCE TRAITS

The impact of stocking density on the growth performance of growing rabbits is listed in Table 2. The results revealed that rabbits stocked at 16 and 24 rabbits/m<sup>2</sup> had lower BW and BWG than those stocked at 8 rabbits/m<sup>2</sup> ( $P < 0.001$ ), with the lowest values were observed in the group of 24 rabbits/m<sup>2</sup>. Also, Rabbits stocked at 16 and 24 rabbits/m<sup>2</sup> exhibited lower FI ( $P = 0.009$ ), and a higher FCR and mortality rate ( $P = 0.001$  and  $0.008$ , respectively).

**Table 2:** Impact of stocking density on growth performance of NZW rabbits.

Items	Stocking density (rabbits/m <sup>2</sup> )			SEM	P value
	8	16	24		
Initial BW, g/ rabbit	571.21	553.45	560.67	10.67	0.593
Final BW, g/ rabbit	2017.83 <sup>a</sup>	1748.67 <sup>b</sup>	1567.58 <sup>c</sup>	20.98	<0.001
BWG, g/rabbit	1446.62 <sup>a</sup>	1195.22 <sup>b</sup>	1006.91 <sup>c</sup>	16.98	<0.001
TFI, g/rabbit	5107.41 <sup>a</sup>	5040.36 <sup>a</sup>	4807.47 <sup>b</sup>	39.43	0.009
FCR (feed: gain)	3.53 <sup>c</sup>	4.22 <sup>b</sup>	4.77 <sup>a</sup>	0.08	0.001
Mortality rate, %	0.00 <sup>c</sup>	5.00 <sup>b</sup>	10.00 <sup>a</sup>	0.13	0.008

Means bearing different superscripts within the same row were significantly different ( $P < 0.05$ ). NZW: New Zealand White rabbits; BW: body weight; TFI: total feed intake; BWG: body weight gain; FCR: feed conversion rate. SEM: standard error of the mean.

### IMPACT OF STOCKING DENSITY ON CARCASS CHARACTERISTICS

Details concerning the features of the carcass are shown in Table 3. No statistical variations ( $P > 0.05$ ) were found between various stocking densities concerning the percentages of skin, full stomach, full intestine, head, kidney, and carcass parts (fore, mid, and hind parts). However, a significant difference ( $P < 0.001$ ) was noticed concerning dressing out and liver percentage, as the rabbits stocked at 24 rabbits/m<sup>2</sup> had a lower dressing and liver percentage as compared with those stocked at 8 and 16 rabbits/m<sup>2</sup>. The SW, CCW, and RCW were significantly decreased as the stocking density increased ( $P < 0.001$ ).

### IMPACT OF STOCKING DENSITY ON HEMATOLOGICAL PARAMETERS

As illustrated in Table 4, the hematological values of RBCs count, Hb, and PCV % were significantly ( $P < 0.01$ ) higher in rabbits stocked at 8 and 16 rabbits/m<sup>2</sup> and lower in

rabbits stocked at 24 rabbits/m<sup>2</sup>. No statistical variations were detected among various stocking densities concerning MCV, MCH, WBCs, lymphocytes, heterophils and H: L ratio.

**Table 3:** Impact of stocking density on carcass traits of NZW rabbits.

Items	Stocking density (rabbits/m <sup>2</sup> )			SEM	P value
	8	16	24		
Slaughter weight (SW), g	2085.71 <sup>a</sup>	1816.21 <sup>b</sup>	1609.07 <sup>c</sup>	25.76	<0.001
Skin, % SW	16.75	17.43	16.89	1.46	0.123
Full stomach, % SW	4.87	5.28	5.67	1.04	0.084
Full intestine, % SW	11.54	12.28	11.93	1.87	0.231
Chilled carcass weight (CCW), g	1202.87 <sup>a</sup>	1034.79 <sup>b</sup>	843.56 <sup>c</sup>	20.87	0.003
Dressing out, %	57.67 <sup>a</sup>	56.97 <sup>a</sup>	52.42 <sup>b</sup>	1.19	<0.001
Head, % CCW	9.52	8.87	9.13	0.88	0.255
Liver, %CCW	6.08 <sup>a</sup>	5.93 <sup>a</sup>	5.12 <sup>b</sup>	0.08	<0.001
Kidney, % CCW	0.99	0.86	0.81	0.43	0.376
Reference carcass weight (RCW), g	997.67 <sup>a</sup>	848.82 <sup>b</sup>	639.56 <sup>c</sup>	16.76	<0.001
Perirenal fat, % RCW	1.45	1.18	1.12	0.67	0.476
Fore part, % RCW	38.23	37.61	37.50	1.73	0.765
Mid part, % RCW	20.87	19.92	19.64	1.68	0.201
Hind part, % RCW	39.27	38.71	38.50	0.93	0.345

Means bearing different superscripts within the same row were significantly different ( $P < 0.05$ ). NZW: New Zealand White rabbits; SEM: standard error of the mean.

**Table 4:** Impact of stocking density on blood hematology of NZW rabbits.

Items	Stocking density (rabbits/m <sup>2</sup> )			SEM	P value
	8	16	24		
RBC, 10 <sup>6</sup> /μl	5.45 <sup>a</sup>	5.27 <sup>a</sup>	4.33 <sup>b</sup>	0.09	0.006
Hb, g/dl	12.45 <sup>a</sup>	11.75 <sup>ab</sup>	10.60 <sup>b</sup>	0.27	0.001
PCV, %	35.35 <sup>a</sup>	34.74 <sup>a</sup>	32.35 <sup>b</sup>	0.36	0.005
MCV, fl	64.35	63.12	62.90	1.65	0.414
MCH, pg	23.91	23.31	23.35	0.87	0.724
WBC, 10 <sup>3</sup> /μl	7.50	8.28	8.74	0.59	0.238
Lymphocytes, %	69.85	70.13	70.25	1.09	0.576
Heterophils, %	36.60	37.51	37.88	0.65	0.167
H: L ratio	0.52	0.53	0.54	0.08	0.379

Means bearing different superscripts within the same row were significantly different ( $P < 0.05$ ). NZW: New Zealand White rabbits; RBC: red blood cell count; Hb: hemoglobin; PCV: packed cell volume; MCV: mean corpuscular volume; MCH: Mean corpuscular hemoglobin; WBC: white blood cell count. SEM: standard error of the mean.

**IMPACT OF STOCKING DENSITY ON SERUM BIOCHEMICAL PARAMETERS**

Data regarding serum biochemical parameters are presented in Table 5. A significant reduction in total protein and globulin concentrations ( $P < 0.001$ ) was reported as the stocking density increased. The concentrations of urea and creatinine were significantly lower ( $P = 0.034$  and  $0.017$ , respectively) in the group having 24 rabbits/m<sup>2</sup> compared with those of 8 and 14 rabbits/m<sup>2</sup>. However, the levels of albumin, cholesterol, triglyceride, AST, and ALT were not significantly influenced by cage density ( $P > 0.05$ ).

**Table 5:** Impact of stocking density on serum biochemical parameters of NZW rabbits.

Items	Stocking density (rabbits/m <sup>2</sup> )			SEM	P value
	8	16	24		
Total protein, g/dl	5.91 <sup>a</sup>	5.21 <sup>b</sup>	4.43 <sup>c</sup>	0.09	<0.001
Albumin, g/dl	3.70	3.53	3.27	0.29	0.089
Globulin, g/dl	2.21 <sup>a</sup>	1.68 <sup>b</sup>	1.16 <sup>c</sup>	0.07	<0.001
Total cholesterol (mg/dl)	78.41	79.50	77.05	2.78	0.895
Triglycerides, mg/dl	126.03	123.50	121.70	4.34	0.536
Urea, mg/dl	31.86 <sup>a</sup>	30.74 <sup>a</sup>	26.36 <sup>b</sup>	1.14	0.034
Creatinine, mg/dl	1.36 <sup>a</sup>	1.25 <sup>a</sup>	0.97 <sup>b</sup>	0.05	0.017
AST, U/l	25.50	24.11	24.67	1.31	0.876
ALT, U/l	81.50	79.21	81.98	2.56	0.937

Means bearing different superscripts within the same row were significantly different ( $P < 0.05$ ). NZW: New Zealand White rabbits; AST: aspartate aminotransferase; ALT: alanine aminotransferase; SEM: standard error of the mean.

**IMPACT OF STOCKING DENSITY ON ECONOMIC PARAMETERS**

Economic calculations revealed that there was no significant difference among various stocking densities concerning feed cost and total costs ( $P > 0.05$ , Table 6). The best values of total and net revenue were observed in rabbits stocked at 8 rabbits/m<sup>2</sup> and the lowest values were recorded in rabbits stocked at 24 rabbits/m<sup>2</sup>. The cost-benefit ratio was significantly increased ( $P < 0.001$ ) from 0.61 to 0.77 as the number of rabbits increased from 8 to 24 rabbits/m<sup>2</sup>.

Stocking density significantly influenced the growth rate of rabbits, with those in high-density groups (16 and 24 rabbits/m<sup>2</sup>) had lower body weight and feed intake and the poorest FCR relative to the low-density group (8 rabbits/m<sup>2</sup>). The lighter body weights that were recorded for rabbits stocked at higher density may be attributable to the lower feed consumption, the lower feed utilization, and the stress that the rabbits exposed due to overcrowding. Similarly, to the current study, it has been reported that high stocking

density decreases body weight and feed consumption and increases FCR in growing rabbits (Mousa-Balabel, 2009; Baiomy, 2012; Trocino et al., 2015). Notably, Amber et al. (2018) reported that the final BW and average FI of ARPI-line rabbits decreased when the density increase from 2 to 4 and 6 rabbits per cage. On contrary, there were no significant differences among rabbits kept at a density of 1, 3, and 5 rabbits/cage (Onbasilar and Onbasilar, 2007), of 4.16, 8.33, 12.5, and 16.67 rabbits/m<sup>2</sup> (Neto et al., 2007), and of 8.6, 12.9, and 17.2 rabbits/m<sup>2</sup> (Dorra et al., 2013) in final BW, BWG, and FI. Also, Volek et al. (2012) reported that stocking density (4 and 10 rabbits/m<sup>2</sup>) did not affect the FCR of Czech white rabbits.

**Table 6:** Impact of stocking density on economic feasibility of NZW rabbits.

Items	Stocking density (rabbits/m <sup>2</sup> )			SEM	P value
	8	16	24		
Feed cost (LE/rabbit)	25.54	25.20	24.04	1.34	0.081
Total cost (LE/rabbit)	55.54	55.20	54.04	1.33	0.123
Total revenue (LE/rabbit)	90.81 <sup>a</sup>	78.69 <sup>b</sup>	70.54 <sup>c</sup>	1.16	<0.001
Net revenue (LE/rabbit)	35.27 <sup>a</sup>	23.49 <sup>b</sup>	16.50 <sup>c</sup>	0.92	<0.001
Cost-benefit ratio	0.61 <sup>c</sup>	0.71 <sup>b</sup>	0.77 <sup>a</sup>	0.02	<0.001

Means bearing different superscripts within the same row were significantly different ( $P < 0.05$ ). Price of kg diet = 5 L.E; price of kg live body weight = 45 L.E. NZW: New Zealand White rabbits; SEM: standard error of the mean.

Consistent with previous reports El-Samra et al. (2013) and Amber et al. (2018), the obtained data demonstrated that carcass traits were not significantly affected by different stocking densities except for dressing out and liver percentage, where the percentages of dressing out and liver of rabbits stocked at 24 rabbits/ m<sup>2</sup> were lower than those reared at 8 and 16 rabbits/m<sup>2</sup>, likely due to the lower carcass weight of rabbits stocked at 24 rabbits/m<sup>2</sup>. However, Villalobos et al. (2008) noted that the high stocking density of 12, 18, and 24 rabbits/m<sup>2</sup> had no significant effect on carcass weight and dressing out of fattening rabbits. Yakubu and Adua (2010) noted that a stocking density of 10, 14.3, 20, and 25 rabbits/m<sup>2</sup> did not affect dressing percentage, carcass weight, and weight of visceral organs.

Our study also showed that high stocking density (24 rabbits/m<sup>2</sup>) significantly diminished estimations of RBCs, Hb, and PCV but had no impact on other blood components, in harmony with the finding of Yakubu et al. (2008) who noted that rabbits stocked at 10 and 14.3 rabbits/m<sup>2</sup> had higher PCV, Hb, and RBCs values relative to those stocked at 20 and 25 rabbits/m<sup>2</sup>, though WBCs and lymphocyte counts did not differ significantly. In contrast, De la Fuente et al. (2004) stated that RBCs

and PCV values were identical in Spanish Giant rabbits housed at 8 and 12 rabbits/cage. Kalaba (2012) observed that California rabbits stocked at 4 rabbits/m<sup>2</sup> had the highest rate of MCV, MCH, and MCHC relative to those stocked at 8, 12, and 16 rabbits/m<sup>2</sup>, while rabbits stocked at 16 rabbits/m<sup>2</sup> had the highest WBCs compared to those kept at 4, 8, and 12 rabbits/m<sup>2</sup>.

Stocking density was found to have no impact on albumin, cholesterol, triglyceride, AST, and ALT concentrations, while the levels of total protein, globulin, urea, and creatinine were significantly decreased as the stocking density increased. Similarly, Kalaba (2012) reported that stocking density (4, 8, 12, and 16 rabbits/m<sup>2</sup>) did not change the levels of albumin and total lipid in rabbit blood plasma, but total protein, globulin, and creatinine were significantly lower in rabbits kept at higher density (16 rabbits/m<sup>2</sup>) than in those housed at lower cage densities. Onbasilar and Onbasilar (2007) noted that cholesterol and triglyceride levels in rabbit serum were not influenced by cage density (2.4, 7.1, and 11.9 rabbits/m<sup>2</sup>). In contrast, Abd El-Monem et al. (2009) concluded that there was no substantial difference in plasma total protein, globulin, and creatinine levels among rabbits housed at four stocking densities (2, 3, 4, and 5 rabbits/cage). El-Samra et al. (2013) observed that by raising the cage density, serum cholesterol, and triglyceride levels were dramatically reduced.

Economical assessment of the experimental groups revealed a substantial decline in the total and net revenue and an increase in cost-benefit ratio as the stocking density increased. In line with the current findings, Amber et al. (2018) noted that as the number of rabbits increased from 2 to 6 per cage, economic efficiency decreased from 1.17 to 0.96, and net income decreased from 18.721 to 13.204 LE. However, Verspecht et al. (2011) showed that decreasing stocking density means recalculating all costs for a decreased number of broiler rabbits, as decreasing stocking density from the standard position of 15 rabbits per m<sup>2</sup> to 10 rabbits per m<sup>2</sup> decreased added value by €22 per doe. Farm income was less and amounted to only €28.10 per doe for the reference situation of 10 rabbits per m<sup>2</sup>.

## CONCLUSIONS AND RECOMMENDATIONS

The results of this study showed that increasing stocking density up to 24 rabbits/m<sup>2</sup> adversely affected economically significant traits and induced stress effects in terms of decreasing protein metabolism (total protein, globulin, and creatinine levels) and some blood disorders.

## AUTHOR'S CONTRIBUTION

MAEO designed the study plan, gathered literature, and

compiled the manuscript. ME assisted in the execution of research work. FAMH provided technical assistance in writing and revising the manuscript. All the authors read and approved the final manuscript.

## CONFLICT OF INTEREST

The authors have declared no conflict of interest.

## REFERENCES

- Abd El-Monem UM, Mahrose KM, Khalil BA (2009). Effect of cage density and climatic condition on the performance of growing rabbits. *Zag. Vet. J.*, 37: 198-208.
- Amber Kh, Eid YZ, Azoz AA, El-Gebaly MM (2018). Effect of stocking density and dietary phytobiotics on growth performance in rabbit during summer season. *J. Sustain. Agri. Sci.*, 44: 13-20.
- Baiomy AA (2012). Growth and carcass traits of New Zealand white rabbits rearing in Upper Egypt as affected by cage density. *Poult. Sci.*, 32: 475-481.
- Bigler L, Oester H (1996). Group housing for male rabbits In: *Proc. 6<sup>th</sup> World Rabbit Congress, Toulouse, France, Vol. 2: 411-415.*
- Dacie JV, Lewis SM (1991). *Practical haematology*, Churchill Livingstone: Edinburgh; New York.
- De la Fuente J, Salazar MI, Ibanez M, Gonzalez de Chavarri E (2004). Effect of season and stocking density during transport on live weight and biochemical measurements of stress, dehydration and injury of rabbits at time of slaughter. *Anim. Sci.*, 78: 285-292. <https://doi.org/10.1017/S1357729800054072>
- Dorra TMI, Ismail FSA, El-Sherif K, Marwa MH (2013). Growth performance of fattening rabbits as affected by stocking density and added dietary organic acids. *J. Anim. Poult. Prod. Mansoura Univ.*, 4: 249-262. <https://doi.org/10.21608/jappmu.2013.71337>
- Egbo ML, Doma UD, Lacdaks AB (2001). Characteristics of small scale rabbit production and management in Bauchi metropolis. *Proc. 26<sup>th</sup> Ann. Conf. Niger. Soc. Anim. Prod., ABU Zaria*, pp. 160-162.
- El-Bayoumi KM, El-Tarabany MS, Abdel-Hamid TM, Sherief WRJA (2018). Effect of stocking density on growth performance, carcass traits and some hematological parameters of New Zealand White rabbits. *Glob. Vet.*, 20: 01-05.
- El-Samra HA, Kalaba ZM, Lamiaa YY (2013). Effect of rearing stocking density on growth performance and some blood parameters of growing rabbits. *J. Anim. Poult. Prod. Mansoura Univ.*, 4: 285-295. <https://doi.org/10.21608/jappmu.2013.71225>
- European Food and Safety Authority (2005). Scientific opinion of the scientific panel on animal health and welfare on "the impact of the current housing and husbandry systems on the health and welfare of farmed domestic rabbit", EFSA-Q-2004-023. *EFSA J.*, 267: 1-31. <https://doi.org/10.2903/j.efsa.2005.267>
- Gondret F, Larzul C, Combes S, de Rochambeau H (2005). Carcass composition, bone mechanical properties, and meat quality traits in relation to growth rate in rabbits. *J. Anim. Sci.*, 83: 1526-1535. <https://doi.org/10.2527/2005.8371526x>
- Grace TIE, Olorunju SAS (2005). Effect of stocking density on performance of growing rabbits in the semi-humid tropics. *Tropicultura*, 23: 19-23.
- Kalaba ZM (2012). Physiological response and stress indicators of California rabbits under intensive conditions in Egypt. *Asian J. Poult. Sci.*, 6: 65-78. <https://doi.org/10.3923/ajpsaj.2012.65.78>
- Mousa-Balabel TM (2009). Effect of cage versus pen and stocking density on behavior and performance in rabbits. *Online J. Vet. Res.*, 13: 48-55.
- National Research Council (1977). *Nutrient requirements of rabbits*, 2nd revised ed., National Academy Press, Washington, DC, USA.
- Neto AC, Lui JF, Dourado LRB, Zanato JAF, Filho EP, Garrafoli NC, da Silva LPG (2007). Effect of stocking density on the performance of growing rabbits. *Biotemas*, 20: 75-79.
- Official methods of analysis (2000). *Association of Official Analytical Chemist*. 17th ed, Virginia, USA.
- Onbasilar EE, Onbasilar I (2007). Effect of cage density and sex on growth, food utilization and some stress parameters of young rabbits. *Scand. J. Lab. Anim. Sci.*, 34: 189-195.
- Samkol P, Lukefahr SD (2008). A challenging role for organic rabbit production toward poverty alleviation in South East Asia. *9<sup>th</sup> World Rabbit Congress, Verona, Italy.*
- Szendrő Zs, Dalle Zotte A (2011). Effect of housing conditions on production and behaviour of growing meat rabbits: A review. *Livest. Sci.*, 137: 296-303. <https://doi.org/10.1016/j.livsci.2010.11.012>
- Trocino A, Filiou E, Tazzoli M, Birolo M, Zuffellato A, Xiccato G (2015). Effect of floor type, stocking density, slaughter age and gender on productive and qualitative traits of rabbits reared in collective pens. *Behav. Welfare Health*, 9: 855-861. <https://doi.org/10.1017/S1751731114003188>
- Verspecht A, Maertens L, Tuytens F, Van Huylenbroeck G, Verbeke W (2011). Economic impact of decreasing stocking densities in broiler rabbit production based on Belgian farm data. *World Rabbit Sci.*, 19: 123-132. <https://doi.org/10.4995/wrs.2011.849>
- Villalobos O, Guillèn O, Garcia J (2008). Effect of cage density on performance of fattening rabbits under heat stress. *Proc. 9<sup>th</sup> World Rabbit Cong. Verona, Italy*, pp. 1631-1639.
- Volek Z, Chodova D, Tumova E, Volkova L, Kudrnova E, Marounek M (2012). Effect of stocking density on growth performance, meat quality and fiber properties of biceps femoris muscle of slow growing rabbits. *Proc. 10<sup>th</sup> World Rabbit Cong. Sharm El-Sheikh, Egypt*, pp. 891-895.
- Yakubu A, Adua MM (2010). Effect of stocking density on growth performance, carcass qualities and cost-benefit of weaned rabbits in the Savanna zone of Nigeria. *Afr. J. Anim. Biomed. Sci.*, 5: 106-109.
- Yakubu A, Adua MM, Adamude H (2008). Welfare and haematological indices of weaner rabbits as affected by stocking density. *Proc. 9<sup>th</sup> World Rabbit Cong. Verona, Italy*, pp. 1269-1273.